In the claims:

Claims 1-19. **(Canceled)**

1 ~~20.~~ **(Previously presented)** A method for treating follicular lymphoma in a subject comprising administering an amount of a composition comprising a soluble lymphotoxin-beta receptor (LT-beta-R) and a pharmaceutically acceptable carrier, such that treatment occurs.

Claims 21-23 **(Canceled)**

2 ~~24.~~ **(Previously presented)** The method of claim ~~20~~ 1 wherein the subject is a mammal.

3 ~~25.~~ **(Previously presented)** The method of claim ~~24~~ 2 wherein the subject is a human.

4 ~~26.~~ **(Previously presented)** The method according to claim ~~20~~ 1 wherein the soluble lymphotoxin-β receptor comprises a ligand binding domain that can selectively bind to a surface LT ligand.

Claims 27-30. **(Canceled)**

5 ~~31.~~ **(Currently amended)** The method of claim ~~20~~ 1, further comprising administering ~~the administration~~ to said subject ~~of~~ at least one chemotherapeutic agent.

6 ~~32.~~ **(Currently amended)** The method of claim ~~20~~ 1, further comprising administering ~~the administration~~ to said subject a ~~of~~ radiation treatments.

7 ~~33.~~ **(Currently amended)** The method of claim ~~20~~ 1 further comprising administering ~~the administration~~ to said subject a ~~of~~ radiation treatments or bone marrow transplant~~ation~~.

Claims 34-35. **(Canceled)**

36. **(Previously presented)** The method of claim 20, wherein the treatment is tumor regression or arrest.

37. **(Previously presented)** The method of claim 20, wherein the soluble LT-beta-R comprises a soluble LT-beta-R fused to one or more heterologous protein domains.

38. **(Currently amended)** The method of claim 37, wherein the soluble LT-beta-R is fused to ~~heterologous protein domain comprises~~ a human immunoglobulin Fc domain.

39. **(Previously presented)** The method of claim 20, wherein the soluble LT-beta-R comprises an extracellular domain of LT-beta-R.

40. **(Previously presented)** The method of claim 20, wherein the soluble LT-beta-R is human LT-beta-R.

41. **(Currently amended)** The method of claim 40, wherein the soluble LT-beta-R ~~immunoglobulin fusion~~ further comprises a human immunoglobulin Fc domain.

42. **(Previously presented)** The method of claim 41, wherein the immunoglobulin is IgG1.

43. **(Previously presented)** A method for disrupting interaction of a B cell lymphoma with its environment in a subject, comprising administering to the subject a composition comprising a soluble LT-beta-R and a pharmaceutically acceptable carrier, such that disruption of the interaction of the B cell lymphoma with its environment occurs.

44. **(Previously presented)** The method of claim 43, wherein the interaction is between the B cell lymphoma and a follicular dendritic cell in the subject.

45. **(Previously presented)** The method of claim 43, wherein the disruption of the interaction results in inhibition of growth of the B cell lymphoma.

18 15
~~46.~~ **(Previously presented)** The method of claim ~~43~~, wherein the soluble LT-beta-R comprises a soluble LT-beta-R fused to one or more heterologous protein domains.

19 18
~~47.~~ **(Previously presented)** The method of claim ~~46~~, wherein the heterologous protein domain comprises a human immunoglobulin Fc domain.

20 15
~~48.~~ **(Previously presented)** The method of claim ~~43~~, wherein the soluble LT-beta-R comprises an extracellular domain of LT-beta-R.

21 15
~~49.~~ **(Previously presented)** The method of claim ~~43~~, wherein the soluble LT-beta-R is human LT-beta-R.

22 21
~~50.~~ **(Previously presented)** The method of claim ~~49~~, wherein the LT-beta-R-immunoglobulin fusion comprises a human immunoglobulin Fc domain.

23 21
~~51.~~ **(Previously presented)** The method of claim ~~49~~, wherein the immunoglobulin is IgG1.

24
~~52.~~ **(New)** A method for treating follicular lymphoma in a human subject, the method comprising administering to the subject a pharmaceutical composition comprising a polypeptide that comprises a soluble, ligand-binding domain of human lymphotoxin-beta receptor (LT-beta-R) fused to a human IgG1 Fc domain, such that treatment occurs.

25
~~53.~~ **(New)** The method of claim 52, wherein the soluble, ligand-binding domain of human LT-beta-R comprises SEQ ID NO:1.